

12038

Feldspathic Basalt

746 grams



Figure 1: Picture of surface of 12038,7 and ,205 showing zap pits. Cube is 1 inch. NASA #S75-34403.

## Introduction

Lunar basalt 12038 is distinctly different from the other Apollo 12 basalts. It has more feldspar, higher Na<sub>2</sub>O and Al<sub>2</sub>O<sub>3</sub>, higher REE and is characterized by a distinct isotopic composition (Nyquist et al. 1981).

According to Hörz and Hartung (1971), rock 12038 has micrometeorite craters on only one surface orientation with a well-developed “soil line” made up of glass splashes and welded dust indicating only one orientation during its history on the lunar surface.

## Petrography

The petrology of 12038 is discussed by Keil et al. (1971), Simpson and Bowie (1971) and Beatty et al. (1979). Beatty et al. describe the texture of 12038 as “hypidiomorphic and dominantly equigranular (~0.6 mm). Plagioclase laths form a loose, randomly oriented network in which pyroxene is either interstitial or partially enclosed. Also present is acicular ilmenite, interstitial cristobalite, trace Ca-phosphate, fayalite, ulvöspinel, K-feldspar, troilite, K-glass and Fe-metal.”




Figure 2: Thin section 12038,29 showing abundant feldspar. NASA photo. Scale is 2 cm.

---

#### Mineralogical Mode of 12038

	Beaty et al. 1979	Neal et al. 1994	Christie et al. 1971
Olivine		0.1	
Pyroxene	49	48.8	55
Plagioclase	44	43.8	30
Ilmenite	3.46	3.5	10
Chromite	0.2	0.2	
“silica”	2.7	2.7	
mesostasis	0.05	0.7	



Figure 3: Photomicrographs of thin section 12038,2 (top = plane-polarized light; bottom = crossed-nicols). Scale 2.2 mm. NASA S70-49437-438.

12038 has an average grain size of about 600 microns (Beatty et al.) and also contains a few % vesicles.

The origin of 12038 has been modeled by Nyquist et al. (1981) in the context of other Apollo 12 basalts. It points to a few % plagioclase in the source region for 12038 magma.

### Mineralogy

**Olivine:** A minor amount of “resorbed” or “cumulus” olivine ( $\text{Fo}_{60}$ ) as well as some fayalite is reported by Keil et al. (1971).

**Pyroxene:** Keil et al. (1971), Busche et al. (1971) and Beatty et al. (1979) reported the pyroxene composition (figure 4). Gose et al. (1972) determined the cell dimensions of exsolved pyroxenes in 12038.




Figure 4: Pyroxene composition for 12038 (adapted from Keil et al. 1971 and Beatty et al. 1977).




Figure 5: Ni and Co in 12038 and 12004 as determined by Simpson and Bowie (1971).

**Pyroxferroite:** Busche et al. (1971) reported pyroxferroite in 12038. Beatty et al. determined that the composition was  $\text{Wo}_{17}\text{En}_3\text{Fs}_{80}$ .

**Plagioclase:** The plagioclase in 12038 ( $\text{An}_{82}$ ) has high Fe content (up to 1.9 wt. %  $\text{FeO}$ , Keil et al. 1971). Beatty et al. (1979) found most plagioclase to be lath-like, typically twinned and zoned from  $\text{An}_{82-73}$ . Christie et al. (1971) observed laths-shaped plagioclase crystals ranging up to 3 mm long occur locally in radiating clusters (figure 3). The last plagioclase to crystallize is anhedral and poikilitically surrounds the earlier formed minerals.

**Ilmenite:** Busche et al. (1971) and Beatty et al. (1979) determined the composition of ilmenite.

**Metallic iron:** Simpson and Bowie (1971) reported low Ni in iron in 12038 (figure 5).




Figure 6: Normalized rare-earth-element diagram for 12038 comparing neutron activation analysis (Brunfelt et al. 1971, Haskin et al. 1971) and spark source mass spectroscopy (Taylor et al. 1971) with isotope dilution mass spectroscopy (line, Schnetzler and Phillipotts 1971, Nyquist et al. 1979).

## Chemistry

12038 has more sodium and aluminum than other Apollo 12 basalts (table 1). The REE pattern is also distinctly different (figure 6). Note the importance of high-quality analyses (and/or correct sample size) is clearly indicated by the dispersion of the data in figure 6.

## Radiogenic age dating

Compston et al. (1971) dated 12038 at  $3.28 \pm 0.21$  b.y. by Rb/Sr (figure 8). Nyquist et al. (1981) obtained ages  $3.35 \pm 0.09$  b.y. by Rb/Sr (figure 9) and  $3.28 \pm 0.23$  b.y. by Nd/Sm (figure 10).

Nyquist showed that the initial Sr and Nd isotope ratios were distinctly different from those of other Apollo 12 basalts (figure 11).

## Cosmogenic isotopes and exposure ages

Burnett et al. (1975) determined an exposure age of  $230 \pm 15$  m.y. by  $^{81}\text{Kr}/^{83}\text{Kr}$ . The suntan age for 12038 (from etched solar flare track studies) is 1.3 m.y. (Bhandari et al. 1971).




Figure 7: Composition of 12038 compared with that of other lunar basalts.

## Other Studies

Biggar et al. (1971) used the composition of 12038 in their experimental studies. At 1 atm. they found olivine was the liquidus phase (1160 C), followed closely by pigeonite and plagioclase, such that they could say 12038 was “multiply saturated” at low pressure.

**Table 1a. Chemical composition of 12038.**

reference weight	LSPET70	Kushiro 71	Cuttita 71	Compston71	Biggar71 Peck	Willis72	Beatty79	Taylor71 ave.
SiO <sub>2</sub> %	49	47.05	(a) 47.1	(b) 46.56	(e)	47.13	(a) 46.61	(e) 46.85
TiO <sub>2</sub>	3.2	3.33	(a) 3.17	(b) 3.31	(e)	3.28	(a) 3.25	(e) 3.25
Al <sub>2</sub> O <sub>3</sub>	12	12.12	(a) 12.8	(b) 12.53	(e)	13.03	(a) 12.45	(e) 12.7
FeO	17	17.91	(a) 17.4	(b) 17.99	(e)	17.73	(a) 17.75	(e) 17.72
MnO	0.26	0.24	(a) 0.24	(b) 0.27	(e)	0.25	(a) 0.251	(e) 0.25
MgO	6.5	7.09	(a) 6.8	(b) 6.71	(e)	6.6	(a) 6.83	(e) 6.74
CaO	11	11.46	(a) 11.4	(b) 11.62	(e)	11.43	(a) 11.48	(e) 11.48
Na <sub>2</sub> O	0.6	0.64	(a) 0.64	(b) 0.66	(e)	0.69	(a) 0.67	(e) 0.67
K <sub>2</sub> O	0.057	0.07	(a) 0.07	(b) 0.073	(e)	0.06	(a) 0.067	(e) 0.07
P <sub>2</sub> O <sub>5</sub>				0.14	(e)	0.14	(a) 0.12	(e) 0.14
S %			(a)		0.06	(e)		0.07 (e) 0.07
<i>sum</i>								
Sc ppm	55		50	(b)			48.2	50 (f)
V	70		126	(b) 104	(e)			120 (f)
Cr	2200	2326	(a) 2210	(b) 1840	(e)	2190	(e) 2050	2100 (f)
Co	23		34	(b) 25	(e)		29.1	29 (f)
Ni	14		6.7	(b) 2	(e)			1 (f)
Cu			10	(b) 8	(e)			5 (f)
Zn				3	(e)			
Ga				3.8	(e)			
Ge ppb								
As								
Se								
Rb	0.7			0.48	(e) 0.6	(d)		
Sr	230		158	(b) 185.8	(e) 186	(d)	185	(e) 173
Y	68		71	(b) 46	(e)		50.5	(e) 59
Zr	260		186	(b) 160	(e)		182	(e) 182
Nb				7	(e)		9.3	(e) 9
Mo								7.8 (f)
Ru								0.05 (f)
Rh								
Pd ppb								
Ag ppb								
Cd ppb								
In ppb								
Sn ppb								100 (f)
Sb ppb								
Te ppb								
Cs ppm							0.02	(f)
Ba	230		142	(b) 120	(e)	107	(e) 119	125 (f)
La				10	(e)		12.47	14 (f)
Ce				25	(e)		30	36 (f)
Pr								5.3 (f)
Nd								26 (f)
Sm								9.4 (f)
Eu							2.2	2.6 (f)
Gd								13 (f)
Tb							1.9	2.9 (f)
Dy								15 (f)
Ho								3.9 (f)
Er								9.9 (f)
Tm								1.6 (f)
Yb		6.3	(b)				6.05	9 (f)
Lu							0.75	
Hf							6.5	6.5 (f)
Ta								
W ppb								
Re ppb								
Os ppb								
Ir ppb								
Pt ppb								
Au ppb								
Th ppm							1	(f)
U ppm							0.25	(f)

technique (a) conventional wet, (b) mixed microchem. XRF emiss. Spec. (c) INAA, (d) IDMS, (e) XRF, (f) SSMS

**Table 1b. Chemical composition of 12038.**

reference weight	Schnetzer71	Brunffelt71	Haskin71	Anders71	Tatsumoto71	Baedecker71	Nyquist79
SiO <sub>2</sub> %							
TiO <sub>2</sub>	3.18	3.4	(h)				
Al <sub>2</sub> O <sub>3</sub>	12.6	12.6	(h)				
FeO	18.14	17.7	(h)				
MnO	0.24	0.32	(h)				
MgO							
CaO							
Na <sub>2</sub> O		0.63	0.63	(h)			
K <sub>2</sub> O	0.076	(d)	0.064	0.6	(h)		0.074 (d)
P <sub>2</sub> O <sub>5</sub>							
S %							
sum							
Sc ppm		44.6	42.7	(h)			
V		154	144	(h)			
Cr		2040	2010	(h)			
Co		28.4	28	(h)	25	(g)	
Ni							
Cu		16.6	10.3	(h)			
Zn		6.1	3.2	(h)	2.09	(g)	1.3 (g)
Ga		5.1	5.5	(h)	4.3	(g)	5.2 (g)
Ge ppb							
As		0.15	0.1	(h)			
Se		0.182	0.174	(h)	0.12	(g)	
Rb	0.604	(d)	0.43	0.44	(h)	0.41	(g)
Sr	190	(d)	137	152	(h)		0.578 (d)
Y							
Zr							
Nb							
Mo							
Ru							
Rh							
Pd ppb							
Ag ppb		80	60	(h)			
Cd ppb					80	(g)	
In ppb		171	149	(h)	5.2	(g)	6.1 (g)
Sn ppb					77	(g)	1.1 (g)
Sb ppb		10	20	(h)			
Te ppb							
Cs ppm		0.014	0.021	(h)	0.015	(g)	
Ba	130	(d)	91	89	(h)		118 (d)
La		11.6	11.5	(h)	11.8	(c)	12.1 (d)
Ce	35	(d)	24	30	(h)	29.1	(c)
Pr							31.7 (d)
Nd	26.3	(d)		22	(c)		23.7 (d)
Sm	8.02	(d)	6.93	6.87	(h)	7.57	(c)
Eu	2.19	(d)	2.04	2.05	(h)	1.969	(c)
Gd	10.6	(d)			10.1	(c)	9.28 (d)
Tb		1.19	1.19	(h)	1.61	(c)	
Dy	11.1	(d)	8.8	8	(h)	9.73	(c)
Ho		1.48	1.48	(h)	1.99	(c)	10.2 (d)
Er	6.12	(d)	5	4.9	(h)	5	(c)
Tm							5.71 (d)
Yb	5.26	(d)	4.9	4.5	(h)	4.8	(c)
Lu	0.814	(d)	0.75	0.76	(h)	0.689	(c)
Hf		5	4.8	(h)			4.64 (d)
Ta		0.59	0.63	(h)			0.65 (d)
W ppb		108	106	(h)			
Re ppb							
Os ppb							
Ir ppb		0.1		(h)			
Pt ppb					0.04	(g)	0.09 (g)
Au ppb		42	3.5	(h)			
Th ppm		0.29	0.18	(h)	0.36	(g)	
U ppm		0.14	0.1	(h)		0.615 (d)	
						0.157 (d)	

technique: (c) INAA, (d) IDMS, (g) RNAA, (h) various NAA




Figure 8: Rb-Sr mineral isochron for 12038 (from Compston et al. 1971).




Figure 9: Rb-Sr mineral isochron diagram for feldspathic basalt 12038 (from Nyquist et al. 1981).

Bogard et al. (1971) reported the content and isotopic composition of rare gases in 12038.




Figure 10: Nd-Sm mineral isochron for 12038 (from Nyquist et al. 1981).




Figure 11: Initial isotopic compositions of 12038 compared with those of Apollo 12 and Luna 24 basalts (Nyquist et al. 1981).

### Processing

A slab (,6) was cut through the middle of 12038 (figure 12) and a column (,8) was cut from the slab (figure 13). A piece of the end (,74) is on public display (figure 14).

### Summary of Age Data for 12038


	Ar/Ar	Rb/Sr	Nd/Sm
Nyquist et al. 1981		$3.35 \pm 0.09$ b.y.	$3.28 \pm 0.23$
Compston et al. 1971		$3.28 \pm 0.21$	

Nyquist 1977 (recalculated)

( $3.22 \pm 0.21$ )

# THE CUTTING OF LUNAR ROCK NO. 12038

DRWG. COMPLETED 6/16/70






Figure 12: Group photo of 12038 after initial cutting showing slab "B" (12038,6) and butt ends 12038,7 and 12038,5 (top facing up). NASA #S70-48268. Scale in cm.






Figure 13: Group photo of column cut from 12038,6 (slab "B"). Scale in cm. NASA #S70-48291.

# THE CUTTING AND CHIPPING OF SLICE 'B'

DRNG. COMPLETED 6/19/70  
(12038.6)







*Figure 14: Lunar basalt 12038, 74 in hermetically-sealed, glass display case.*